

An ionic liquid of the formula K^+A^- wherein K^+ and A^- are as defined herein, are suitable for use in electrochemical cells and capacitors. These ionic liquids can also be used in catalysis, as inert solvents, and as hydraulic liquids.

bioRxiv preprint doi: https://doi.org/10.1101/076891; this version posted November 1, 2016. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted bioRxiv a license to display the preprint in perpetuity. It is made available under aCC-BY-NC-ND 4.0 International license.	
Figure S1	
A	
<p>Number of reads</p>	
B	
<p>Number of reads</p>	
C	
<p>Number of reads</p>	
D	
<p>Number of reads</p>	
E	
<p>Number of reads</p>	
F	
<p>Number of reads</p>	
G	
<p>Number of reads</p>	
H	
<p>Number of reads</p>	
I	
<p>Number of reads</p>	
J	
<p>Number of reads</p>	
K	
<p>Number of reads</p>	
L	
<p>Number of reads</p>	
M	
<p>Number of reads</p>	
N	
<p>Number of reads</p>	
O	
<p>Number of reads</p>	
P	
<p>Number of reads</p>	
Q	
<p>Number of reads</p>	
R	
<p>Number of reads</p>	
S	
<p>Number of reads</p>	
T	
<p>Number of reads</p>	
U	
<p>Number of reads</p>	
V	
<p>Number of reads</p>	
W	
<p>Number of reads</p>	
X	
<p>Number of reads</p>	
Y	
<p>Number of reads</p>	
Z	
<p>Number of reads</p>	